
AD-A037 105

GEORGETOWN UNIV WASHINGTON D C
THERMODYNAMICS AND STATISTICAL MECHANICS OF LIQUID CRYSTALS AND-ETC(U)
1977

UNCLASSIFIED

ARO-6425.18-C

END
DATE
PLANCE
PLAN



Unclassified SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered) READ INSTRUCTIONS REPORT DOCUMENTATION PAGE BEFORE COMPLETING FORM 2. GOVT ACCESSION ENT'S CATALOG NUMBER 6425.18-C Final Report THERMODYNAMICS AND STATISTICAL MECHANICS OF Sep 66 - 31 Dec 76 LIQUID CRYSTALS AND THEIR SOLUTIONS. Daniel E. Martire DA-ARO-D-31-124-G8175 DA-ARO-D-31-124-G1167 DA-ARO-D-31-124-73-G2 9. PERFORMING ORGANIZATION NAME AND ADDRESS Georgetown University Washington, D. C. 20007 11. CONTROLLING OFFICE NAME AND ADDRESS 1977 U. S. Army Research Office Post Office Box 12211 Research Triangle Park NC 27700
MONITORING IGENCY NAME & ADDRESS(If different from Controlling Office) 15. SECURITY CLASS. (of this report) Unclassified DECLASSIFICATION DOWN GRADING 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 17. DISTRIBUTION STATEMENT (of the obstract entered in Block 20, if different from Report) 18. SUPPLEMENTARY NOTES The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents. 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Thermodynamic properties Solutions Liquid crystals Solvents Phase transformations Mixtures Statistical mechanics Binary systems (materials)


ABSTRACT (Continue on reverse side if necessary and identify by block number)

This is a brief summary of the liquid crystal research supported by ARO at Georgetown University during the past ten years. Research has been directed toward a more detailed understanding, on a molecular level, of the thermodynamic properties of (a) phase transitions in liquid crystals and (b) binary mixtures of nonmesomorphic solutes and nematic (or cholesteric) solvents.

FINAL REPORT

- 1. ARO PROJECT NUMBER: CRD-AA-IP-6425-C
- 2. PERIOD COVERED BY REPORT: 1 September 1966 31 December 1976*
- 3. TITLE OF PROJECT: Thermodynamics and Statistical Mechanics of Liquid Crystals and Their Solutions
- 4. GRANT NUMBER: DA-ARO-D-31-124-G817 (9/1/66 8/31/69)
 DA-ARO-D-31-124-G1167 (9/1/69 8/31/72)
 DA-ARO-D-31-124-73-G2 (9/1/72 12/31/76)*
- 5. NAME OF INSTITUTION: Georgetown University
- 6. AUTHOR OF REPORT: Daniel E. Martire, Professor
- 7. LIST OF MANUSCRIPTS PUBLISHED UNDER ARO SPONSORSHIP DURING THIS PERIOD, INCLUDING JOURNAL REFERENCES: Listed following Summary of Research
- 8. SCIENTIFIC PERSONNEL SUPPORTED (6 MONTHS OR MORE) BY THIS PROJECT AND DEGREES AWARDED DURING THIS REPORTING PERIOD:
 - (a) Daniel E. Martire, Principal Investigator
 - (b) David G. Willey, Postdoctoral Research Associate
 - (c) Göran I. Agren, Postdoctoral Research Associate
 - (d) George M. Janini, Postdoctoral Research Associate
 - (e) Martha A. Cotter, Graduate Student (Ph.D., with distinction, June 1969)
 - (f) Laurence C. Chow, Graduate Student (Ph.D., June 1970)
 - (g) Joel M. Schnur, Graduate STudent (Ph.D., January 1972)
 - (h) Henry T. Peterson, Graduate Student (Ph.D., with distinction, January 1973)
 - (i) Gwo-Chung Lin, Graduate Student (Ph.D., June 1975)
 - (j) Ghassan A. Oweimreen (Ph.D. expected June 1977)

Performance period of the grant extended by 16 months (9/1/75 - 12/31/76) without additional funding.



Daniel E. Martine January 14, 1977

SUMMARY OF RESEARCH

The liquid crystal research supported by ARO at Georgetown University during these past ten years has been directed toward a more detailed understanding, on a molecular level, of the thermodynamic properties of (a) phase transitions in liquid crystals and (b) binary mixtures of nonmesomorphic solutes and nematic (or cholesteric) solvents.

The potential of gas-liquid chromatography (glc) for studying the thermodynamics of infinitely dilute solutions of nonmesomorphic solutes in liquid-crystalline solvents was demonstrated in an exploratory investigation (1). Subsequent experimental studies confirmed the validity of our glc approach by establishing the absence of surface effects (3,10) and the agreement between glc data and extrapolated static measurements (11). The glc method and a statistical-thermodynamic model were then successfully employed in obtaining and interpreting thermodynamic solution quantities for a wide range of solutes in nematic and cholesteric solvents (8,13,15,22). Our understanding of the unique solvent properties of liquid crystals has been advanced through these studies. Other novel applications of glc were developed: analytical separation of geometric isomers using a nematic liquid phase (4); determination of nematic order parameters by a solute probe method (9); and complementary thermodynamic studies to interpret the effect of organic solutes on the selective reflectance of visible light by cholesteric mixtures (12). Apparatus were designed and tested (11,20) for the thermodynamic study of mixtures of nonmesopmorphic solutes at finite concentration in nematic solvents. Experiments were conducted to investigate the effect of solute size, shape and flexibility on nematic phase stability (14,20).

Several statistical-mechanical models were developed to provide the necessary framework for interpreting the thermodynamic properties of liquid-crystalline phase transitions in one and two component systems. A scaled-particle theory (5,6,7) of rigid rods (one component) demonstrated the primary role of repulsive forces (or packing effects) in the nematic-isotropic liquid transition. A lattice model (2) of rigid rods was used to investigate the secondary effect of attractive forces. The results of both models suggested the probable importance of endchain flexibility in homologous series. Accordingly, a rotationalisomeric model (15) and a lattice model of particles consisting of rigid central cores and semiflexible pendant groups (17) were developed to investigate the effect of end-chain flexibility. The results of these models (and available thermodynamic data) strongly indicated that the nematic-isotropic transition can be adequately described in terms of molecular packing and flexibility effects, with attractive forces providing a uniform background and having only a minor effect on the transition properties. A simple tunnel model of rigid rods (one component) was developed and successfully applied to interpret the thermodynamic behavior of the smectic A - nematic phase transition (21). Finally, models involving two components - mixtures of rigid rods of different lengths (16) and mixtures of rigid rods and spheres (18,19,20) were developed to aid in interpreting the experimental results (14,20) for solute-induced nematic-isotropic phase transitions.

LIST OF MANUSCRIPTS

- Thermodynamics of Solutions with Liquid Crystal Solvents. I. A GLC Study of Cholesteryl Myristate, Journal of Physical Chemistry, 72, 3489 (1968). (with P.A. Blasco, P.F. Carone, L.C. Chow and H. Vicini).
- A Quasi-Chemical Lattice Treatment of Rod-like Molecules. Application to the Nematic-Isotropic Transition in Liquid Crystals, Molecular Crystals and Liquid Crystals, 7, 295 (1969). (with M.A. Cotter).
- Thermodynamics of Solutions with Liquid Crystal Solvents. II. Surface Effects with Nematogenic Compounds, Journal of Physical Chemistry, 73, 1127 (1969). (with L.C. Chow).
- 4. Gas Chromatography and Thermodynamics of Divinylbenzene Separations on 4,4'-Dihexoxyazoxybenzene Liquid Crystal, Analytical Chemistry, 42, 176 (1970). (with W.L. Zielinski, Jr., D.H. Freeman and L.C. Chow).
- 5. Statistical Mechanics of Rod-like Particles. I. A Scaled Particle Treatment of a Fluid of Perfectly Aligned Rigid Cylinders, Journal of Chemical Physics, 52, 1902 (1970). (with M.A. Cotter).
- 6. Statistical Mechanics of Rod-like Particles. II. A Scaled Particle Investigation of the Aligned+Isotropic Transition in a Fluid of Rigid Spherocylinders, <u>Journal of Chemical Physics</u>, <u>52</u>, 1909 (1970). (with M.A. Cotter).
- 7. Statistical Mechanics of Rod-like Particles. III. A Scaled Particle Treatment of a Fluid of Rigid Spherocylinders with Restrictired Orientational Freedom, <u>Journal of Chemical Physics</u>, <u>53</u>, 4500 (1970). (with M.A. Cotter).
- 8. Thermodynamics of Solutions with Liquid Crystal Solvents. III. Molecular Interpretation of Solubility in Nematogenic Solvents, Journal of Physical Chemistry, 75, 2005 (1971). (with L.C. Chow).
- Thermodynamics of Solutions with Liquid Crystal Solvents. IV. Gas-Liquid Chromatography Determination of the Degree of Order in the Nematic Mesophase, <u>Molecular Crystals and Liquid Crystals</u>, <u>14</u>, 293 (1971). (with L.C. Chow).
- 10. Thermodynamics of Solutions with Liquid Crystal Solvents. V. Surface Effects with Cholesteric Compounds, Analytical Chemistry, 43, 1201 (1971). (with J.M. Schnur).
- 11. Activity Coefficient of n-Heptane in 4,4'-Di-hexoxyazoxybenzene Liquid Crystal, Journal of Physical Chemistry, 76, 596 (1972). (with H.T. Peterson and W. Lindner).
- 12. Thermodynamics of Solutions with Liquid Crystal Solvents. VI. The Effect of Organic Solutes on the Selective Reflectance of Visible Light by Cholesteric Mixtures, Molecular Crystals and Liquid Crystals, 18, 55 (1972). (with D.G. Willey).

- 13. Thermodynamics of Solutions with Liquid Crystal Solvents. VII.

 A Mixture of Cholesteryl Chloride and Cholesteryl Myristate,

 Molecular Crystals and Liquid Crystals, 26, 213 (1974). (with

 J.M. Schnur).
- 14. Thermodynamics of Solutions with Liquid Crystal Solvents. VIII. Solute Induced Nematic-Isotropic Transitions, Molecular Crystals and Liquid Crystals, 25, 89 (1974). (with H.T. Peterson).
- 15. On the Question of Molecular Flexibility in Nematogenic Compounds, Molecular Crystals and Liquid Crystals, 28, 63 (1974).
- 16. Lattice Model for a Binary Mixture of Hard Rods of Different Lengths. Application to Solute Induced Nematic-Isotropic Transitions, <u>Journal of Chemical Physics</u>, 61, 3547 (1974). (with H.T. Peterson and M.A. Cotter).
- 17. End-Chain Flexibility and the Nematic-Isotropic Transition in Liquid Crystals: A Lattice Model of Hard Particles with Rigid, Rodlike, Central Cores and Semiflexible Pendant Segments, Journal of Chemical Physics, 61, 3959 (1974). (with G.I. Agren).
- 18. Lattice Model For a Binary Mixture of Hard Rods and Hard Cubes. Application to Solute Induced Nematic-Isotropic Transitions, Journal de Physique, 36, Cl-141 (1975). (with G.I. Agren).
- 19. Mixture of Hard Spherocylinders and Spheres in the Virial Expansion, Physical Review A, 11, 1040 (1975). (with G.I. Agren).
- 20. The Effect of Quasispherical Solutes on the Nematic to Isotropic Transition in Liquid Crystals, Journal of Chemical Physics, 64, 1456 (1976). (with G.A. Oweimreen, G.I. Agren, S.G. Ryan and H.T. Peterson).
- 21. A Tunnel Model for the Nematic to Smectic A Transition in Liquid Crystals, <u>Journal of Chemical Physics</u>, in preparation. (with G.C. Lin and L.C. Chow).
- 22. Thermodynamics of Solutions with Liquid Crystal Solvents. IX. Non-mesomorphic Solutes in Nematic and Isotropic MBBA, Journal of Physical Chemistry, in preparation. (with G.C. Lin and G.A. Oweimreen).

Note:

- a. D.E. Martire is one of the authors of each of the above manuscripts.
- b. Reprints of manuscripts number 21 and 22, which are largely based on G.C. Lin's doctoral dissertation (see following page for abstract), will be sent to ARO following publication of the papers.

Thermotropic Liquid Crystals: A Cell Model for the Smectic A-Nematic Transition and A Thermodynamic Investigation of Dilute Solutions in Nematic and Isotropic MBBA (Ph.D. Dissertation of Gwo-Chung Lin)

Abstract

In order to gain a better understanding of the fundamental nature of the smectic A-nematic transition in liquid crystals, a simple cell model is applied to a fluid of aligned, rigid, cylinders in Part I. Particle motion is separated into components parallel and perpendicular to the long axis of the molecules. The partition functions are formulated by considering the contribution from particles localized within their cells and the contribution from positional disorder in the nematic phase. The transition temperature is determined by equating the Helmholtz free energies of the two phases, in the limit of zero external pressure.

When realistic values for the dimensional and energetic parameters are used in the model calculations, reasonable values are obtained for the transition temperature, absolute density and configurational energy. It is found that: (a) the smectic A phase becomes more favored (i.e., the transition temperature increases) as the strength of segmental core-core interactions and/or the length of the central core increases, (b) the transition enthalpy, entropy and fractional volume change decrease with increasing terminal interactions (ϵ '), while the transition temperature is virtually insensitive to ϵ '. The most significant success of the model is its ability to produce the observed trends in the transition temperature and entropy in homologous series.

In Part II, gas-liquid chromatography is used to obtain infinite dilution activity coefficients and partial molar enthalpies and entropies of solution for 26 nonmesomorphic solutes in both the nematic and isotropic phases of N-(p-methoxy-benzylidene)-p-n-butylaniline (MBBA). The data are interpreted in the light of Chow and Martire's, and Humphries, James and Luckhurst's statistical thermodynamic solution models, and are compared with results obtained for other nematogenic compounds.



